



Only an island as impudent as Cephalonia would have the insouciance to situate itself upon a faultline that exposes it to the recurrent danger of cataclysmic earthquakes.

Captain Corelli's Mandolin (1994)

Louis de Bernières


Argostoli Bay, Sunday, October 5, 2025

Cephalonia: A story of a natural laboratory for seismic hazard studies since 2011 and a reference site SIGMA3 studies



Fabrice HOLLENDER (CEA, ISTERre), Nikolaos THEODOULIDIS (ITS AK), Pierre-Yves BARD (ISTERre), Armand MARSICAL (ISTERre), Agisilaos KONIDARIS (Ionian University), Zafeiria ROUMELIOTI (Univ. Patras), Kiriaki KONSTANTINIDOU (ITS AK), Pauline RISCHETTE (CEA, ISTERre), Vincent PERRON (CEA, ISTERre), Paola TRAVERSA (EdF), Elias EI HABER (EdF) and many others...



The island of Cephalonia, the town of Argostoli, the ARGONET vertical array



Seismotectonic context



The 1953 earthquake

- 7.2 on August 12, 1953, at 9:24 GMT
- 450 to 600 deaths
- After the EQ 100,000 inhabitants out of a total population of approximately 125,000 left the island.




The first seismological studies carried out on Cephalonia

Early seismological measurements: Ionianet

Period: July 1998 – October 2000

61 events, 3 stations → 182 records

$3.5 \leq M \leq 5.3$ $10 \text{ km} \leq R \leq 125 \text{ km}$



2011: The European NERA program



ROMA meeting, 2-4 March 2011
Minutes of sub-meeting on Data Sharing Scheme and New Experiment
Roma, INGV, March 3, 2011

Participants for Data Sharing Scheme & New Experiment

-
AUTH: K. Ptilakis (first part then moved to modelers)
INGV: G. Cultrera, G. Milana, G. DiGiulio, F. Cara, D. Famiani, P. Bordononi
SED: D. Faeh, Jan Burjanek
ITSAK: N. Theodoulidis
GFZ: S. Paolai, Ang. Strollo
ISTERRE: C. Cornou

- Proposed for new experiment for surface topography effects (more geophysical work needed)
- Selected broadband Italian stations for surface topography effects
 - Geodata available
 - Waveforms available
- Cavola site – landslide
 - Geodata available
 - Waveforms available
- Grenoble area – basin effects
 - Geodata available
 - Waveforms available
- Grevena – basin effects
 - Geodata available
 - Waveforms available
- Argostoli – basin effect
 - Geodata available
 - Waveforms available
 - **Proposed for new experiment for basin and surface topography effects***

NERA program experiments



2011: The NERA program

Bull Earthquake Eng (2018) 16:529–560
https://doi.org/10.1007/s10518-017-0225-4

ORIGINAL RESEARCH PAPER

Basin effects on ground motion: the case of a high-resolution experiment in Cephalonia (Greece)

N. Theodoulidis¹ · G. Cultrera² · C. Cornou³ · P.-Y. Bard⁴ · T. Boxberger⁵ · G. DiGiulio⁶ · A. I. D. Kementzidou^{1,6} · K. Makra⁷ · The Argostoli NERA Team

Received: 13 February 2017 / Accepted: 3 September 2017 / Published online: 15 February 2018
© Springer Science+Business Media B.V. 2017

Abstract In this study data and results of a high-resolution experiment in Cephalonia (Greece) regarding empirical basin effects are presented. Accelerometers were deployed in the basin of Argostoli for 7 months (September 2011–April 2012). Due to high seismicity in the surrounding area this array recorded thousands of local earthquakes. Data used in this work come from a selection of 162 earthquakes with $3 \text{ km} \leq R \leq 600 \text{ km}$, with magnitude range, $1.0 \leq M \leq 4.5$.

Argostoli NERA Team: Ch. Androu (Technological Educational Institute of Cephalonia, Greece), R. Baur (GFZ German Research Centre for Geosciences, Helmholtz-Centre, Potsdam, Germany), S. Bayle (Institut des Sciences de la Terre (ISTerre), Université de Grenoble, France), G. DiGiulio (GFZ German Research Centre for Geosciences, Helmholtz-Centre, Potsdam, Germany), G. Cultrera (Istituto Nazionale di Geofisica e Vulcanologia (INGV), Roma, Italy), R. Cogliati (Istituto Nazionale di Geofisica e Vulcanologia (INGV), Roma, Italy), C. Cornou (Institut des Sciences de la Terre (ISTerre), Université de Grenoble, France), A. Fodarella (Istituto Nazionale di Geofisica e Vulcanologia (INGV), Roma, Italy), E. Günther (GFZ German Research Centre for Geosciences, Helmholtz-Centre, Potsdam, Germany), A. Konidaris (Technological Educational Institute of Ionian Islands, Argostoli, Greece), Institut des Sciences de la Terre (ISTerre), Université de Grenoble, France, German Research Centre for Geosciences, Helmholtz-Centre, Potsdam, Germany, Research Centre for Geosciences, Helmholtz-Centre, Potsdam, Germany), Geofisica e Vulcanologia (INGV), Roma, Italy), G. Riccio (Istituto Nazionale di Geofisica e Vulcanologia (INGV), Roma, Italy), and A. Savvaidis (Institute of Engineering Seismology and Earthquake Engineering (ITSAK), Thessaloniki, Greece).

✉ N. Theodoulidis
nthoo@tsak.gr

G. Cultrera
giuliana.cultrera@ingv.it

C. Cornou
cecile.cornou@univ-grenoble-alpes.fr

P.-Y. Bard
Pierre-Yves.Bard@univ-grenoble-alpes.fr

T. Boxberger
tobias.boxberger@gfz-potsdam.de

**ARGOSTOLI EXPERIMENT
[16/9/2011 - 16/4/2012]**

...and AFTER WHAT?

NERA Meeting in Rome, 26-28 March, 2012

multidimensional site effects

Julien P. France

Surface geology and geometry are known to be important for the amplification of surface waves. As a consequence, understanding of physical patterns is required to characterize and quantify them. However, the complexity of such effects as well as the variability in case of small sedimentary valleys. The propagating across two 2-D dense seismic basins to the extent to which they are contributing to the ground motion. The array was deployed in the small-size, shallow alluvial basin of Argostoli, Greece, and consisted of three stations ranging from 5 to 160 m. A set of 40 stations at epicentral distances up to 200 km, was used to characterize the seismicity. The phase velocities crossing the array were extracted, and Rayleigh waves was obtained. The results show that surface waves (up to 60 per cent of total energy), with a dominant frequency (~ 1.5 Hz) of the valley (~ 3 Hz) while Rayleigh waves dominate in a given frequency range. The outcomes of this research provide a method to study the 2-D/3-D site effects in Argostoli. The method applied here is useful for evaluating seismic ground motion.

Keywords: Site effects; Wave propagation; Seismicity

Surface geology and geometry are known to be important for the amplification of locally generated surface waves (e.g. Kennen 1996). Especially, surface waves significantly modify the wavefield on, leading to large amplifications, significant ground strains (Moczo & Bard 2007; Bard et al. 1999; Cornou et al. 2003a,b; Pini & Pini 2010). Such motions can be dangerous on structures and lifelines, particularly in sedimentary valleys where edge-induced waves reverberate within shallow layers and contribute to the generation

Nikos Theodoulidis EPPO-ITSAK


The SINAPS@ age




SINAPS@ project

- March 11, 2011: Accident of Fukushima
- Call for projects from the French agency ANR on topics related to nuclear safety
- SINAPS@ project accepted, including the deployment of a vertical accelerometric array in Argostoli (= Argonet)
- September 2013: first geophysical survey on Cephalonia Island
- January 26, 2014: M6.1 earthquake in Argostoli
- January 27-28, 2024: Official Kick-off meeting of SINAPS@, decision to launch a post-seismic survey

The January 26, 2014 earthquake



Peak Ground Acceleration Map USGS
 ShakeMap: 1km ENE of Lixourion, Greece
 Jan 26, 2014 13:55:42 UTC M6.1 N38.21 E20.45 Depth: 8.0km ID:usb000m8ch



PGA (%g)	0.1	0.2	0.5	1	2	5	10	20	50	100	200
----------	-----	-----	-----	---	---	---	----	----	----	-----	-----


Scale based on Worden et al. (2012)

Version 1: Processed 2020-06-06T08:06:04Z

△ Seismic Instrument ○ Reported Intensity



★ Epicenter

The SINAPS@ post-seismic team experienced the second earthquake!



The February 6, 2014 earthquake

Fabrice's awakening
Vincent's awakening
Um... in your opinion, what magnitude is it?
Awareness that this is a major earthquake...
We leave the hotel (without taking the elevator...)
We end the night in the car.



Post-seismic survey



26/1/14 earthquake



03/2/14 earthquake




Difficult fieldwork to meet scientific objectives



During the equipment test (Monday, February 3), an average of one replica was measured every 30 seconds.

Post-seismic survey

1075 aftershocks on February 12, 2014




During the equipment test (Monday, February 3), an average of one replica was measured every 30 seconds.





Scientific objectives of the mission

Several thousand earthquakes recorded




Assessment of site effects in the Koutavos basin



Perron et al., 2018


Scientific objectives of the mission

Evaluation of the spatial coherence of seismic motion on a rock and sediment network





Svay et al., 2017
Imtiaz et al., 2018

Measurement of the rotational component of seismic motion




Sbaa et al., 2017

3D geological model




Cushing et al., 2020

The ARGONET vertical accelerometric array









4 borehole accelerometers at different depths
 2 surface accelerometers (sediment & rock)
 2 surface rotation sensors
 2 meteorological stations



Installed in July 2015,
 Database (version 2025) : more than 1500 events with an
 excellent signal-to-noise ratio. Open data.



Measurement highlighting seasonal variations and slight non-linearities



The early SIGMA-3 age



Building effect of seismic recordings

West side view




■ STA : in the basement with LEF2

East side view




- Pauline Rischette's thesis
- 3 sites over 9 on Cephalonia




Topography effect: ArgoScarp experiment



Topography effect: ArgoScarp experiment





Topography effect: ArgoScarp experiment



- **Highest SSRs** reached in the **longitudinal** direction of the cliff
- **Lowest SSRs** reached in the **transversal** direction of the cliff

ArgoSlab



ArgoDAS-1 and dense node array



- Optical fiber
- ▲ 3C seismometers (nodes)

The SIGMA-3 next age




October 2025 campaign: a very broad partnership!




Rémy BURLLOT (for CEA), Eirini CHATZIANAGNOSTOU (CEA/ISTerre), Cécile CORNOU (ISTerre), Frédéric DADOUN (Geodesia), Elias El HABER (EdF), Fabrice HOLLENDER (CEA), Philippe REIFFSTECK (UGE), Zafeiria ROUMELIOTI (Univ. Patras), Nikos THEODOULIDIS (ITS AK), Paola TRAVERSA (EdF), drillers from GEOT.ER, 'trencher' team from Tzouganatos, etc.

The BENTO benchmark

- BENTO – BENCHMARK of 3D Numerical simulations on TOPographic sites
- Organized as part of the upcoming ESG2026 conference, in collaboration with CEA–EdF–ISTerre
- Phase 1: verification on “canonical” cases.
- Phase 2: validation on a small area with very strong short-wavelength curvature (ArgoScarp data).
- Phase 3: validation on a larger area with long-wavelength curvature (measurements on Mount Ainos).



ArgoScarp experiment, BENTO phase 3



**Expérience ArgoScarp :
effets topo à courte longueur d'onde**

**Base de données utilisée pour le benchmark
"BENTO"**

Launch of measurements for BENTO Phase 3

- 18 measurement points for a duration of 3 to 4 months



Launch of measurements for BENTO Phase 3



Launch of measurements for BENTO Phase 3



Launch of measurements for BENTO Phase 3



'Only an island as indolent as Cephalonia would allow itself to be infested by such troupes of casual and impertinent goats'

Captain Corelli's Mandolin (1994),

Louis de Bernières

Two jurists preparing a collaboration agreement and discussing intellectual property

Drilling operation



- Installation of optical fiber on boreholes
- Installation of pore pressure probes
- In situ non-linearity parameter with cyclic test

Installation of pore-pressure sensors

- 4 sensors at 3.5, 5.5, 7.5 and 15.5 m (zone of low CPT N-values, accelerometer depth)
- Vibrating-wire cone-tip sensors (GeoSense probe), to be pushed to the bottom of the borehole




Installation of pore-pressure sensors




Installation of pore-pressure sensors



First earthquake recorded with the pressure sensors




Location: 6 Km WNW from Assos Kefallinia
Magnitude: 3.0 ML
Origin Time: 12/10/2025 11:37:39.73 (GMT)
Latitude: 38.4004 N
Longitude: 20.4836 E
Focal Depth: 13.1 Km
Type: Manual



ARGO-DAS2


- DAS (Distributed Acoustic Sensing):
 - Principle: the optical fiber acts as a dense array of sensors, continuously measuring strain variations along its path using laser backscattering.
 - Interest: enables the acquisition of seismic measurements with very high spatial resolution (metric scale) over several kilometers, without deploying point sensors.
- ARGODAS1 (2023): initial measurements taken as part of a collaboration between CEA and ISTERre (but measurement issues with the interrogator worked only 9 days of data were collected).
- ARGODAS2 (2025): new measurements on the same fiber, but with the addition of fibers in boreholes.
 - Approximately 1,400 m of horizontal fiber.
 - + 2 boreholes (20 and 50 m) of vertical fiber.
 - + 62 nodes for validation/calibration of DAS measurements.
 - Collaboration between EdF, CEA, and ISTERre as part of SIGMA-3.
 - Fibers and nodes installed, DAS measurements themselves over a period of 2 to 3 months (starting in November)



ARGO-DAS2



ARGO-DAS2




In situ cyclic pressure tests




- Collaboration between EdF, CEA, and Gustave Eiffel University as part of SIGMA-3:
- 2 x 18 measurements (from 3 to 20 m) of nonlinear parameters using a cyclic pressure gauge

In situ cyclic pressure tests



ARGONET, current instrumentation setup



Scientific production from Sinaps@ and Sigma programs from Cephalonia experiments: 15 peer-reviewed articles'



Cushing EM, Hollender F, Moiriat D, Guyonnet-Benaize C, Theodoulidis N, Pons-Branchu E, S  p  lcre S, Bard P-Y, Cornou C, Dechamp A, Mariscal A, Roumelioti Z (2020) Building a three dimensional model of the active Plio-Quaternary basin of Argostoli (Cephalonia Island, Greece): An integrated geophysical and geological approach. *Engineering Geology* 265:105441. <https://doi.org/10.1016/j.enggeo.2019.105441>

Grendas I, Theodoulidis N, Bard P-Y, Perron V, Hatzidimitriou P, Hollender F (2022a) Can site effects be estimated with respect to a distant reference station? Performance of the spectral factorization of coda waves. *Geophysical Journal International* 230:1–28. <https://doi.org/10.1093/gji/ggac040>

Grendas I, Theodoulidis N, Hollender F, Hatzidimitriou P (2022b) Spectral decomposition of S-waves in investigating regional dependent attenuation and improving site amplification factors: A case study in western Greece. *Bull Earthquake Eng* 20:6441–6465. <https://doi.org/10.1007/s10518-022-01459-z>

Grendas I, Hollender F, Perron V, Theodoulidis N, Buscetti M, Traversa P (2025a) The SSRc method as a practical solution for evaluating site response using far apart stations: applications in Western Greece and South-Eastern France. *Bull Earthquake Eng*. <https://doi.org/10.1007/s10518-025-02221-x>

Grendas I, Roumelioti Z, Theodoulidis N, Hollender F (2025b) Ground-Motion Dependency on Seasonal Variations: Observations at the ARGONET Array, Cephalonia, Greece. *Bulletin of the Seismological Society of America* 115:1801–1817. <https://doi.org/10.1785/0120240175>

Hollender F, Roumelioti Z, Maufroy E, Traversa P, Mariscal A (2020) Can We Trust High-Frequency Content in Strong-Motion Database Signals? Impact of Housing, Coupling, and Installation Depth of Seismic Sensors. *Seismological Research Letters*. <https://doi.org/10.1785/0220190163>

Imtiaz A, Perron V, Hollender F, Bard P, Cornou C, Svay A, Theodoulidis N (2018) Wavefield Characteristics and Spatial Incoherency: A Comparative Study from Argostoli Rock- and Soil-Site Dense Seismic Arrays. *Bulletin of the Seismological Society of America* 108:2839–2853. <https://doi.org/10.1785/0120180025>

Perron V, G  lis C, Froment B, Hollender F, Bard P-Y, Cultrera G, Cushing EM (2018a) Can broad-band earthquake site responses be predicted by the ambient noise spectral ratio? Insight from observations at two sedimentary basins. *Geophysical Journal International* 215:1442–1454. <https://doi.org/10.1093/gji/ggy355>

Perron V, Hollender F, Mariscal A, Theodoulidis N, Andreou C, Bard P, Cornou C, Cottareau R, Cushing EM, Frau A, Hok S, Konidaris A, Langlaude P, Laurendeau A, Savvaids A, Svay A (2018b) Accelerometer, Velocimeter Dense-Array, and Rotation Sensor Datasets from the Sinaps@ Postseismic Survey (Cephalonia 2014–2015 Aftershock Sequence). *Seismological Research Letters* 89:678–687. <https://doi.org/10.1785/0220170125>

Roumelioti Z, Hollender F, Gu  g  n P (2020) Rainfall-Induced Variation of Seismic Waves Velocity in Soil and Implications for Soil Response: What the ARGONET (Cephalonia, Greece) Vertical Array Data Reveal. *Bulletin of the Seismological Society of America* 110:441–451. <https://doi.org/10.1785/0120190183>

Roumelioti Z, Hollender F, Theodoulidis N, Grendas I (2025) Variation in Seismic Wave Velocities at Shallow Depth and the Masking of Nonlinear Soil Behavior Based on the ARGONET (Cephalonia, Greece) Vertical Array Data. *Applied Sciences* 15:10727. <https://doi.org/10.3390/app151910727>

Sbaa S, Hollender F, Perron V, Imtiaz A, Bard P-Y, Mariscal A, Cochard A, Dujardin A (2017) Analysis of rotation sensor data from the SINAPS@ Kefalonia (Greece) post-seismic experiment—link to surface geology and wavefield characteristics. *Earth, Planets and Space* 69. <https://doi.org/10.1186/s40623-017-0711-6>

Svay A, Perron V, Imtiaz A, Zentner I, Cottareau R, Clouteau D, Bard P-Y, Hollender F, Lopez-Caballero F (2017) Spatial coherency analysis of seismic ground motions from a rock site dense array implemented during the Kefalonia 2014 aftershock sequence. *Earthquake Engineering & Structural Dynamics* 46:1895–1917

Theodoulidis N, Hollender F, Mariscal A, Moiriat D, Bard P-Y., Konidaris A, Cushing M, Konstantinidou K, Roumelioti Z (2018) The ARGONET (Greece) Seismic Observatory: An Accelerometric Vertical Array and Its Data. *Seismological Research Letters* 89:1555–1565. <https://doi.org/10.1785/0220180042>

Touhami S, Lopez-Caballero F, Clouteau D (2020) A holistic approach of numerical analysis of the geology effects on ground motion prediction: Argostoli site test. *J Seismol*. <https://doi.org/10.1007/s10950-020-09961-0>



Thank you for you attention!

The major milestones of ARGONET and Cephalonia test-site

